

1. Find allowed energies of the *half* harmonic oscillator

$$V(x) = \begin{cases} \frac{1}{2}m\omega^2 x^2, & x > 0, \\ \infty, & x < 0. \end{cases}$$

2. A charged particle (mass m , charge q) is moving in a simple harmonic potential (frequency $\omega/2\pi$). In addition, an external electric field \mathcal{E}_0 is also present. Write down the hamiltonian of this particle. Find the energy eigenvalues, eigenfunctions. Find the average position of the particle, when it is in one of the stationary states.
3. Assume that the atoms in a CO molecule are held together by a spring. The spacing between the lines of the spectrum of CO molecule is 2170 cm^{-1} . Estimate the spring constant.
4. If the hermite polynomials $H_n(x)$ are defined using the generating function $G(x, s) = \exp(-s^2 + 2xs)$, that is

$$\exp(-s^2 + 2xs) = \sum_n \frac{H_n(x)}{n!} s^n,$$

- (a) Show that the Hermite polynomials obey the differential equation

$$H_n''(x) - 2xH_n'(x) + 2nH_n(x) = 0$$

and the recurrence relation

$$H_{n+1}(x) = 2xH_n(x) - 2nH_{n-1}(x).$$

- (b) Derive Rodrigues' formula

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} e^{-x^2}.$$

5. Let ϕ_n be the n th stationary state of a particle in harmonic oscillator potential. Given that the lowering operator is

$$\hat{a} = \frac{1}{\sqrt{2\hbar m\omega}} (m\omega \hat{X} + i\hat{P}).$$

and $\xi = \sqrt{m\omega/\hbar}x$,

- (a) show that

$$\hat{a} = \frac{1}{\sqrt{2}} \left(\xi + \frac{d}{d\xi} \right).$$

- (b) Show that

$$\hat{a}\phi_n(\xi) = \sqrt{n}\phi_{n-1}(\xi)$$

6. Let $B = \{\phi_n | n = 0, 1, \dots\}$ be the set of energy eigenfunctions of the harmonic oscillator. Find the matrix elements of \hat{X} and \hat{P} wrt to basis B
7. Suppose that a harmonic oscillator is in its n^{th} stationary state.

- (a) Compute uncertainties σ_x and σ_P in position and momentum. [Hint: To calculate expectation values, first write \hat{X} and \hat{P} in terms of the lowering operator \hat{a} and its adjoint.]
- (b) Show that the average kinetic energy is equal to the average potential energy (Virial Theorem).

8. A particle of mass m in the harmonic oscillator potential, starts out at $t = 0$, in the state

$$\Psi(x, 0) = A(1 - 2\xi)^2 e^{-\frac{m\omega}{2\hbar}\xi^2}$$

where A is a constant and $\xi = \sqrt{m\omega/\hbar}x$.

- (a) What is the average value of energy?
- (b) After time T , the wave function is

$$\Psi(x, T) = B(1 + 2\xi)^2 e^{-\frac{m\omega}{2\hbar}\xi^2}$$

for some constant B . What is the smallest value of T ?

9. Let ϕ_n be eigenstates of the harmonic oscillator. For a given complex number μ , let

$$\chi_\mu = e^{-\frac{|\mu|}{2}} \sum_{n=0}^{\infty} \frac{\mu^n}{\sqrt{n!}} \phi_n.$$

Such states are called *coherent* states.

- (a) Show that

$$\hat{a}\chi_\mu = \mu\chi_\mu$$

that is χ_μ is an eigenstate of \hat{a} .

- (b) If the state of the oscillator is χ_μ , then show that $\sigma_x\sigma_p = \hbar/2$.
- (c) The state of the oscillator $\Psi(t = 0) = \chi_\mu$, then show that

$$\Psi(t) = \chi_{\mu'}$$

where $\mu' = e^{-i\omega t}\mu$. That means, if the state of the system, at an instant is a coherent state, then it is a coherent state at all times.

- (d) Optional: If you choose the hilbert space to be $L_2(\mathbb{R})$, then show that $|\Psi(x, t)|^2$ is a gaussian wave packet and the wave packet performs a harmonic oscillations without changing the shape.