

Quadratic approximation


Prof. (Dr.) Rajib Kumar Bhattacharjya



Professor, Department of Civil Engineering
Indian Institute of Technology Guwahati, India

Room No. 005, M Block

Email: rkbc@iitg.ernet.in, Ph. No 2428




$$f(x) = 2x^4 - x^3 + 5x^2 - 12x + 1$$

$$f'(x) = 8x^3 - 3x^2 + 10x - 12 = 0$$

Solving for x

$$x^* = 0.8831 \text{ and } f(x^*) = -5.1702$$



$$f(x) = 2x^4 - x^3 + 5x^2 - 12x + 1$$

Quadratic approximation of the function at x_0 can be written as

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + 0.5*f''(x_0)(x - x_0)^2$$

Approximate function for $x_0 = 0$


Now we can minimize the function

$$\text{Minimize } f'(x_0)(x - x_0) + 0.5*f''(x_0)(x - x_0)^2$$

Solution is

$x^* = 1.2$ and $f(x^*) = -3.7808$ and $f'(x^*) = 9.5040$

This is the solution of the approximate function: First trial



$$f(x) = 2x^4 - x^3 + 5x^2 - 12x + 1$$

Quadratic approximation of the function at x_0 can be written as

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + 0.5 * f''(x_0)(x - x_0)^2$$

Approximate function for $x_0 = 1.2$


Now we can minimize the function

$$\text{Minimize } f'(x_0)(x - x_0) + 0.5 * f''(x_0)(x - x_0)^2$$

Solution is

$$x^* = 0.9456, f(x^*) = -5.1229 \quad \text{and } f'(x^*) = 1.5377$$

This is the solution of the approximate function: Second trial



$$f(x) = 2x^4 - x^3 + 5x^2 - 12x + 1$$

Quadratic approximation of the function at x_0 can be written as

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + 0.5*f''(x_0)(x - x_0)^2$$

Approximate function for $x_0 = 0.9456$


Now we can minimize the function

$$\text{Minimize } f'(x_0)(x - x_0) + 0.5*f''(x_0)(x - x_0)^2$$

Solution is

$x^* = 0.8864$ and $f(x^*) = -5.1701$ and $f'(x^*) = 0.0785$

This is the solution of the approximate function: Third trial



$$f(x) = 2x^4 - x^3 + 5x^2 - 12x + 1$$

Quadratic approximation of the function at x_o can be written as

$$f(x) = f(x_o) + f'(x_o)(x - x_o) + 0.5*f''(x_o)(x - x_o)^2$$

Approximate function for $x_o = 0.8864$

Now we can minimize the function

$$\text{Minimize } f'(x_o)(x - x_o) + 0.5*f''(x_o)(x - x_o)^2$$

Solution is

$$x^* = 0.8831 \text{ and } f(x^*) = -5.1702 \text{ and } f'(x^*) = 0.00099$$



This is the solution of the approximate function: Fourth trial

**STOP
ITERATION**

Gradient is negligible

Now take an example of multivariable problem

$$\text{Minimize } f(x_1, x_2) = (x_1^2 + x_2 - 11)^2 + (x_1 + x_2^2 - 7)^2$$



$$\text{Minimize } f(x_1, x_2) = (x_1^2 + x_2 - 11)^2 + (x_1 + x_2^2 \pm 7)^2 \quad x_o = [2 \ 2]^T$$



The quadratic approximation of the function at $x_o = [2 \ 2]^T$ can be written as

$$f(X) = f(X_o) + (X - X_o)\nabla f(X_o)^T + (X - X_o)H(X_o)(X - X_o)^T$$

For first approximation

$$\text{Minimize} \quad f(X) = (X - X_o)\nabla f(X_o)^T + (X - X_o)H(X_o)(X - X_o)^T$$

$$\text{Or,} \quad f(X) = [x_1 - 2 \quad x_2 - 2] \begin{bmatrix} -42 \\ -18 \end{bmatrix} + [x_1 - 2 \quad x_2 - 2] \begin{bmatrix} 14 & 16 \\ 16 & 30 \end{bmatrix} \begin{bmatrix} x_1 - 2 \\ x_2 - 2 \end{bmatrix}$$




Solution


Trial	X value	Gradient
1	7.9268	-0.5610
2	5.7945	-4.4555
3	4.4415	-3.1670
4	3.7952	-2.3927
5	3.6086	-1.9928
6	3.5858	-1.8623
7	3.5844	-1.8483
8	3.5844	-1.8481
9	3.5844	-1.8481

Optimal solution


Gradient is almost negligible



$$\text{Minimize } f(x) = x^3 + 2 * x^2 + \frac{x}{2} - 1$$



$$x_0 = 0$$



$$x_0 = -2$$

Thanks
R.K. Bhattacharyya CE/ITG