

Assignment 1

1. If f is continuously differentiable on S^1 , then show that $\hat{f}'(n) = in\hat{f}(n)$ for all $n \in \mathbb{Z}$. Deduce that there exists $C > 0$ such that $|\hat{f}(n)| \leq \frac{C}{|n|}$. Does the above conclusion hold if f is absolutely continuous?
2. If f is a function of bounded variation on $[-\pi, \pi]$, then show that $|\hat{f}(n)| \leq \frac{\text{Var}(f)}{2\pi|n|}$ for all $n \in \mathbb{Z}$.
3. For $f \in L^1(S^1)$, show that $\hat{f}(n) = \frac{1}{4\pi} \int_{-\pi}^{\pi} [f(x) - f(x + \frac{\pi}{n})] e^{-inx} dx$. Further, use this identity to prove the Riemann-Lebesgue lemma.
4. Let $f \in L^1(S^1)$ be such that $|f(x+h) - f(x)| \leq M|h|^\alpha$, for some $0 < \alpha < 1$ and $M > 0$ and for all $x, h \in S^1$. Show that $\hat{f}(n) = O\left(\frac{1}{|n|^\alpha}\right)$.
5. Show that the Fejer's kernel F_n can be expressed as $F_n(t) = \sum_{j=-n}^n \left(1 - \frac{|j|}{n}\right) e^{ijt}$.
6. Let $f \in L^1(S^1)$ and $m \in \mathbb{N}$ and define $f_m(t) = f(mt)$. Prove that $\hat{f}_m(n) = \hat{f}\left(\frac{n}{m}\right)$, if $(m, n) \neq 1$ and $\hat{f}_m(n) = 0$ otherwise.
7. Let f be a function on S^1 . For $x, y \in S^1$, define $\tau_x f(y) = f(x - y)$. Show that $x \rightarrow \tau_x f$ is continuous in $L^p(S^1)$ for $1 \leq p < \infty$. That is, $\|\tau_x f - f\|_p \rightarrow 0$ when $|x| \rightarrow 0$. Does the above conclusion hold if $p = \infty$?
8. If $f \in L^1(S^1)$ and $g \in L^\infty(S^1)$, then show that $\lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t)g(nt)dt = \hat{f}(0)\hat{g}(0)$.
9. For given $f \in L^1(S^1)$, define an operator T_f on $L^1(S^1)$ by $T_f(g) = f * g$. Show that T_f is a bounded operator on $L^1(S^1)$ and $\|T_f\| = \|f\|_1$.
10. Let P be a trigonometric polynomial of degree n on S^1 . Show that $\|P'\|_\infty \leq 2n\|P\|_\infty$.
11. Let $1 \leq p \leq \infty$ and $p^{-1} + q^{-1} = 1$. For $f \in L^p(S^1)$ and $g \in L^q(S^1)$, show that $f * g$ is a continuous function on S^1 .
12. Suppose $f \in L^\infty(S^1)$ satisfies $|\hat{f}(n)| \leq \frac{k}{|n|}$ for some $k > 0$ and for all $n \in \mathbb{Z} \setminus \{0\}$. Prove that $|S_n(f)(t)| \leq \|f\|_\infty + 2k$, where $S_n(f) = D_n * f$.
13. If f is a bounded monotone function on S^1 , then show that $\hat{f}(n) = O\left(\frac{1}{|n|}\right)$.
14. If f is Riemann integrable on $[-\pi, \pi]$, then show that $\sum_{n=-\infty}^{\infty} |\hat{f}(n)|^2 < \infty$ and hence $\hat{f}(n) = o(1)$.
15. Prove that, if the series of complex numbers $\sum_{n=0}^{\infty} a_n$ converges to s , then $\sum_{n=0}^{\infty} a_n$ is Cesaro as well as Abel summable to s .
16. Prove that, if the series of complex numbers $\sum_{n=0}^{\infty} a_n$ is Cesaro summable to σ , then $\sum_{n=0}^{\infty} a_n$ is Abel summable to σ . However, converse need not be true.

17. Let $u(r, \theta) = \frac{\partial P_r}{\partial \theta}(\theta)$, where $P_r(\theta)$ is the Poisson kernel defined on the open unit disc $\mathbb{D} = \{re^{i\theta} : 0 \leq r < 1, \theta \in [-\pi, \pi]\}$. Show that $\Delta u = 0$ on \mathbb{D} and $\lim_{r \rightarrow 1} u(r, \theta) = 0$ for all $\theta \in [-\pi, \pi)$.
18. Let f be Riemann integrable on $[-\pi, \pi]$, and $A_r(f)(\theta) = f * P_r(\theta)$ for $0 \leq r < 1$, denotes the Abel mean of f . If f has jump discontinuity at θ , then show that $\lim_{r \rightarrow 1} A_r(f)(\theta) = \frac{f(\theta^+) + f(\theta^-)}{2}$. Justify that $\lim_{r \rightarrow 1} A_r(f)(\theta) \neq \frac{f(\theta)}{2}$, when f is continuous at θ .
19. Let f be Riemann integrable on $[-\pi, \pi]$, and $\sigma_n(f)(\theta) = f * F_n(\theta)$, where F_n is Fejer's kernel. If f has jump discontinuity at θ , then show that $\lim_{n \rightarrow \infty} \sigma_n(f)(\theta) = \frac{f(\theta^+) + f(\theta^-)}{2}$.
20. Let f be Riemann integrable on $[-\pi, \pi]$ such that $\hat{f}(n) = O\left(\frac{1}{|n|}\right)$ for all $n \in \mathbb{Z}$.
- (a) Show that $S_N(f)(\theta) = D_N * f(\theta) \rightarrow f(\theta)$ if f is continuous at θ .
- (b) If f has jump discontinuity at θ , then show that $S_N(f)(\theta) \rightarrow \frac{f(\theta^+) + f(\theta^-)}{2}$.
- (c) If f is continuous on $[-\pi, \pi]$, then $S_N(f) \rightarrow f$ uniformly.
21. For $f \in L^2(S^1)$, show that $\frac{1}{n} \sum_{k=0}^{n-1} f(x + \frac{k}{n}) \rightarrow \hat{f}(0)$ as $n \uparrow \infty$ in the metric of $L^2(S^1)$.
22. Does there exist a function $f \in L^1(S^1)$ such that $\sum_{n=-\infty}^{\infty} |\hat{f}(n)|^2 = \infty$?
23. If $f \in L^1(S^1)$ vanishes near $x = 0$, then show that $S_N(f) \rightarrow 0$ uniformly near $x = 0$.
24. Let f be a function on $[-\pi, \pi]$ such that $|f(\theta) - f(\varphi)| \leq M|\theta - \varphi|$, for some $M > 0$ and for all $\theta, \varphi \in [-\pi, \pi]$.
- (a) For $u(r, \theta) = P_r * f(\theta)$, show that $\frac{\partial u}{\partial \theta}$ exists for all $0 \leq r < 1$ and $|\frac{\partial u}{\partial \theta}| \leq M$ for some $M > 0$.
- (b) Show that $\sum_{n=-\infty}^{\infty} |\hat{f}(n)| \leq |\hat{f}(0)| + 2M \sqrt{\sum_{n=1}^{\infty} \frac{1}{n^2}}$.
25. If f is continuously differentiable on S^1 , then show that $\sum_{n=-\infty}^{\infty} (1 + |n|^2) |\hat{f}(n)|^2 < \infty$.
26. If $\{G_n\}_{n=1}^{\infty}$ is a family of good kernels on S^1 , then show that $\lim_{n \rightarrow \infty} \hat{G}_n(k) = 1$.
27. Let f and g be Riemann integrable on $[-\pi, \pi]$. Define $\tilde{g}(x) = \overline{g(-x)}$.
- (a) Show that $\frac{1}{2\pi} \int_{-\pi}^{\pi} |g(t)|^2 dt = g * \tilde{g}(0)$.
- (b) $\frac{1}{2\pi} \int_{-\pi}^{\pi} |f * g(x)|^2 dx = \frac{1}{2\pi} \int_{-\pi}^{\pi} |f * \tilde{g}(x)|^2 dx$.
28. Let $f \in L^1(S^1)$ be such that $\hat{f}(|n|) = -\hat{f}(-|n|) \geq 0$ for all $n \in \mathbb{Z}$. Show that $\sum_{n>0} \frac{\hat{f}(n)}{n} < \infty$.
29. If $\{K_n\}_{n=1}^{\infty}$ and $\{J_n\}_{n=1}^{\infty}$ are families of good kernels on S^1 , then show that $\{K_n * J_n\}_{n=1}^{\infty}$ is a family of good kernels.
30. Suppose f is an absolutely continuous function on S^1 such that $f' \in L^2(S^1)$. Prove that $\sum_{n=-\infty}^{\infty} |\hat{f}(n)| \leq \|f\|_1 + 2 \sqrt{\sum_{n=1}^{\infty} \frac{1}{n^2}} \|f'\|_2$.